arXiv:chao-dyn/9707001v1 27 Jun 1997

Comment on “Optimal Periodic Orbits of Chaotic Systems”

Scott M. Zoldi*† and Henry S. Greenside*
Department of Physics
Duke University, Durham, NC 27708-0305
(June 25, 1997)

In a recent Letter [1] (see also [2]), Hunt and Ott argued that *short*-period unstable periodic orbits (UPOs) would be the invariant sets associated with a chaotic attractor that are most likely to optimize the time average of some smooth scalar performance function. In this Comment, we show that their conclusion does not hold generally and that optimal time averages may specifically require long-period UPOs. This situation can arise when long-period UPOs are able to spend substantial amounts of time in a region of phase space that is close to large values of the performance function.

Our counterexample is based on a numerical study of the Lorenz equations [3] for the classical parameter values $r = 28$, $\sigma = 10$, and $b = 8/3$, for which most initial conditions evolve towards the familiar butterfly-shaped chaotic attractor. We chose as a physically-motivated performance function the Nusselt number $N(t)$ which is a dimensionless spatially-averaged measure of the vertical heat transport of a convection cell [4]; for the Lorenz equations, it is given in terms of the Lorenz variables x and y by

$$N = 1 + \frac{2}{br} xy. \tag{1}$$

We evaluated the time average of this expression, $\langle N \rangle_t$, for each of 718 distinct UPOs of the Lorenz equations which had been calculated by a damped-Newton numerical method described elsewhere [5]. A comparison of the calculated UPOs with a known binary symbolic dynamics showed that all UPOs of period $T \leq 6.0$ were found by our method.

Our results are summarized in Fig. 1. Panel (a) shows how the time-averaged Nusselt number $\langle N \rangle_t$ depends on the period T of the UPOs. The shortest period UPOs give average heat transport values that are less than the average value for the chaotic attractor itself (indicated by the horizontal line at $N = 2.68$). The largest values of $\langle N \rangle_t$ occur with *increasing* period. All these values should be compared with the still larger value $N = 2.93$ that is attained for the two nonzero unstable fixed points $x = y = \pm\sqrt{b(r-1)}, z = r - 1$. Panel (b) shows that the long-period large-Nusselt-number UPOs achieve a larger value of $\langle N \rangle_t$ specifically because they are asymmetric and spend more time in the vicinity of a nonzero fixed point. Symbolic dynamics predicts the existence of these asymmetric UPOs up to an arbitrarily long period. Since the fixed points lie distinctly outside the Lorenz attractor, the Nusselt number of all UPOs should be strictly bounded by the fixed-point value.



FIG. 1. **(a)** Time-averaged Nusselt number $< N >_t$ versus period T for unstable period orbits (UPOs) associated with the Lorenz attractor. **(b)** Projection on the $x - y$ plane of a long-period UPO with $T = 9.82$ and $N = 2.750$, showing the asymmetry that allows the UPO to spend much time near a fixed point with large Nusselt number.

We thank Brian Hunt and Ed Ott for discussions about the topic of this Comment. This research was supported by a DOE Computational Science Graduate Fellowship, by NSF grants NSF-DMS-93-07893 and NSF-CDA-92123483-04, and by DOE grant DOE-DE-FG05-94ER25214.

---

* Also Center for Nonlinear and Complex Systems, Duke U., Durham, NC
† E-mail: zoldi@phy.duke.edu

[1] B. R. Hunt and E. Ott, Phys. Rev. Lett. **76**, 2254 (1996).
[2] B. R. Hunt and E. Ott, Phys. Rev. E **54**, 328 (1996).
[3] E. N. Lorenz, J. Atmos. Sci **20**, 130 (1963).
[4] F. H. Busse, Rep. Prog. Phys. **41**, 1929 (1978).
[5] S. Zoldi and H. Greenside, submitted to PRL; LANL number chao-dyn/9704005 (unpublished).